

A Bibliography on Body Weight Support Gait Training

Listings in RED indicate citations from the year 2002 & later

† Denotes articles that involve or discuss Body Weight Support Therapy

* Denotes articles that use or are about the LiteGait or another Mobility Research Product

Basic Science

1. REVIEW

- Andersson O, Forssberg H, Grillner S, Wallen P.(1981) Peripheral feedback mechanisms acting on the central pattern generators for locomotion in fish and cat. *Canadian Journal of Physiology and Pharmacology*, 59, 713-726.
- Armstrong D.M. (1986). Supraspinal contributions to the initiation and control of locomotion in the cat. *Progress in Neurobiology*, 26, 273-361.
- Arshavsky YI, Deliagina TG, Orlovsky GN (1997). Pattern Generation. *Current Opinion in Neurobiology*, 7, 781-789.
- Burke RE, Degtyarenko AM, Simon ES. (2001). Patterns of locomotor drive to motoneurons and last-order interneurons: clues to the structure of the CPG. *Journal of Neurophysiology*, 86(1), 447-62.
- Dettmann M, Linder M, Sepic S. (1987) Relationship among walking performance, postural stability and functional assessment of the hemiplegic patient. *Am J Phys Med*, 66, 77-90.
- Duysens J, Clarac F, Cruse H. (2000) Load-regulating mechanisms in gait and posture: Comparative aspects. *Physiological Reviews*, 80, 83-133.
- Hultborn H, et al (1998) How do we approach the locomotor network in the mammalian spinal cord? *Annals of the New York Academy of Sciences*, 860, 70-82.
- Jordan LM. (1998) Initiation of locomotion in mammals. *Annals of the New York Academy of Sciences*, 860, 83-93.
- Mori, S. (1998) Neurophysiology of locomotion: Recent advances in the study of locomotion. *Annals of the New York Academy of Sciences*, 860, 55-78.
- Pearson KG, Misiaszek JE, Fouad K. (1998) Enhancement and resetting of locomotor activity by muscle afferents. *Annals of the New York Academy of Sciences*, 860, 203-215.
- Pearson KG. (1995) Proprioceptive regulation of locomotion. *Current Opinion in Neurobiology*, 5, 786-791.
- Rossignol S. (1996) Neural control of stereotypic limb movements. In: Rowell LB, Shepherd JT. *Handbook of Physiology; Section 12, Exercise: Regulation and integration of multiple systems*. New York: John Wiley. 174-216.
- Taub E, Uswatte G, Pidikiti R. (1999) Constraint-Induced movement therapy: A new family of techniques with broad application to physical rehabilitation- A clinical review. *Journal of Rehabilitation Research and Development*, 35(3), 237-251
- Whelan PJ.(1996) Control of locomotion in the decerebrate cat. *Progress in Neurobiology*, 49, 481-515.

Basic Science

2. LOCOMOTION

- Andersson O, Grillner S. (1983) Peripheral control of the cat's step cycle II. *Acta Physiologica Scandinavica*, 118, 229-239.
- Andersson O, Grillner S. (1981) Peripheral control of the cat's step cycle. I. Phase dependent effects of ramp-movements of the hip during "fictive locomotion". *Acta Physiologica Scandinavica*, 113, 89-101.
- Beresovskii VK. (1990) Structure of spinal cord locomotor strip in the cat. *Acta Neurobiologiae Experimentalis*, 50, 41-46.
- Blaszczyk J, Loeb GE. (1993). Why cats pace on the treadmill. *Physiology and Behavior*, 53, 501-507.
- Borojerdi B, Ferbert A, Foltys H, Kosinski CM, Noth J, Schwarz M. (1999) Evidence for a non-orthostatic

- origin of orthostatic tremor. *Journal of Neurology, Neurosurgery, and Psychiatry*, 66(3), 284-8.
- Canu MH, Falempin M. (1998) Effect of hindlimb unloading on interlimb coordination during treadmill locomotion in the rat. *European Journal of Applied Physiology and Occupational Physiology*, 78(6), 509-15.
- Dale N, Kuenzi F. (1997) Ionic currents, transmitters and models of motor pattern generators. *Current Opinion in Neurobiology*, 7(6), 790-6.
- De Leon RD, London NJS, Roy RR, Edgerton VR. (1999) Failure analysis of stepping in adult spinal cats. *Progress in Brain Research*, 123, 341-348.
- Deliagina TG, Fagerstedt P. (2000) Responses of reticulospinal neurons in intact lamprey to vestibular and visual inputs. *Journal of Neurophysiology*, 83, 864-878.
- Deliagina TG, Zelenin PV, Fagerstedt P, Grillner S, Orlovsky GN.(2000) Activity of reticulospinal neurons during locomotion in the freely behaving lamprey. *Journal of Neurophysiology*, 83, 853-863.
- Deliagina TG, Orlovsky GN, Pavlova GA. (1983). The capacity for generation of rhythmic oscillations is distributed in the lumbosacral spinal cord of the cat. *Experimental Brain Research*. 53, 81-90.
- Dietz V. (1998) Evidence for a load receptor contribution to the control of posture and locomotion. *Neuroscience and Biobehavioral Reviews*, 22, 495-499.
- Drew T, Dubuc R, Rossignol S.(1986) Discharge patterns of reticulospinal and other reticular neurons on chronic, unrestrained cats walking on a treadmill. *Journal of Neurophysiology*, 55, 375-401.
- Duysens J, Van de Crommert HWAA. (1998) Neural control of locomotion; Part 1: The central pattern generator from cats to humans. *Gait and Posture*, 7, 131-141.
- Eidelberg E, Story JL, Walden JG, Meyer BL.(1981) Anatomical correlates of return of locomotor function after partial spinal cord lesions in cats. *Experimental Brain Research* 42(1), 81-88.
- Field EC, Stein PSG. (1997) Spinal cord coordination of hindlimb movements in the Turtle: Intralimb temporal relationships during bilateral scratching and swimming. *Journal of Neurophysiology*, 78, 1394-1403.
- Forsberg H, Grillner S, Halbertsma J, Rossignol S. (1980). The locomotion of the low spinal cat. II. Interlimb coordination. *Acta Physiologica Scandinavica*, 108, 283-295.
- Forsberg H, Grillner S, Halbertsma J. (1980) The locomotion of the low spinal cat. I. Coordination within a hindlimb. *Acta Physiologica Scandinavica*, 108, 269-281.
- Forsberg H, Grillner S, Rossignol S.(1975) Phase dependent reflex reversal during walking in chronic spinal cats. *Brain Research*, 85, 103-107.
- Fukuyama H, et al. (1997). Brain functional activity during gait in normal subjects: A SPECT study. *Neuroscience Letters*, 228, 183-186.
- Grillner S, Ekeberg O, Manira AE, Lansner A, Parker D, Tegner J, Wallen P.(1998) Intrinsic function of a neuronal network – a vertebrate central pattern generator. *Brain Research Review*, 26, 184-197.
- Grillner S, Dubuc, R. (1988) Control of locomotion in vertebrates: Spinal and supraspinal mechanisms. *Advances in Neurology*, 47, 425-453.
- Grillner S, Zangger P. (1984).The effect of dorsal root transection on the efferent motor pattern in the cat's hindlimb during locomotion. *Acta Physiology Scand Scandinavica*, 120, 393-405.
- Kably B, Drew T.(1998) Corticoreticular pathways in the cat. II. Discharge activity of neurons in area 4 during voluntary gait modifications. *Journal of Neurophysiology*, 80(1), 406-24.
- Leblond H, Menard A, Gossard JP. (1998)Vestibulo- and reticulospinal control of the extensor half-center in locomotion. *Annals of the New York Academy of Sciences*, 860, 563-565.
- Mackay M. (2002) Central Pattern Generation of Locomotion: A Review of the Evidence. *Physical Therapy*, 82.1, 69-83.
- Massion J, Dufosse M. (1988) Coordination between posture and movement: why & how? *NIPS*, 3,88-93
- Matsuyama K, Drew T. (2000).Vestibulospinal and reticulospinal neuronal activity during locomotion in the

- intact cat. II. Walking on an inclined plane. *Journal of Neurophysiology*, 84(5), 2257-76.
- Matsuyama K, Drew T. (2000). Vestibulospinal and reticulospinal neuronal activity during locomotion in the intact cat. I. Walking on a level surface. *Journal of Neurophysiology*, 84 (5), 2237-56.
- Mori S, Matsui T, Kuze B, Asanome M, Nakajima K, Matsuyama K. (1999). Stimulation of a restricted region in the midline cerebellar white matter evokes coordinated quadrupedal locomotion in the decerebrate cat. *Journal of Neurophysiology*, 82, 290-300.
- Mori S, Matsui T, Kuze B, Asanome M, Nakajima K, Matsuyama K. (1998) Cerebellar-induced locomotion: Reticulospinal control of spinal rhythm generating mechanism in cats. *Neuroscience Research*, 30, 257-69
- Pearson KG, Rossignol S. (1991). Fictive motor patterns in chronic spinal cats. *Journal of Neurophysiology*, 66, 1874-1887.
- Prentice SD, Drew T. (2001). Contributions of the reticulospinal system to the postural adjustments occurring during voluntary gait modifications. *Journal of Neurophysiology*, 85(2), 679-98.
- Rossignol S, Chau C, Brustein E, Giroux N, Bouyer L, Barbeau H, Reader TA. (1998) Pharmacological activation and modulation of the central pattern generator for locomotion in the cat. *Annals of the New York Academy of Sciences*, 860, 346-359.
- Smith JL, Smith LA, Zernicke RF, Hoy M. (1982) Locomotion in exercised and nonexercised cats cordotomized at two or twelve weeks of age. *Experimental Neurology*, 76, 393-413.
- Stein PSG, Victor JC, Field EC, Currie SN. (1995). Bilateral control of hindlimb scratching in the spinal turtle: Contralateral spinal circuitry contributes to the normal ipsilateral motor pattern of fictive rostral scratching. *Journal of Neuroscience*, 15, 4343-4355.

Basic Science

3. LEARNING

- Barbeau H, Rossignol S. (1987). Recovery of locomotion after chronic spinalization in the adult cat. *Brain Research*, 412, 84-95.
- Brustein E, Rossignol S. (1998) Recovery of locomotion after ventral and ventrolateral spinal lesions in the cat. I. Deficits and adaptive mechanisms. *Journal of Neurophysiology*, 80, 1245-1267.
- De Leon RD, Tamaki H, Hodgson JA, Roy RR, Edgerton VR. (1999) Hindlimb locomotor and postural training modulates glycinergic inhibition in the spinal cord of the adult spinal cat. *Journal of Neurophysiology*, 82, 359-369.
- De Leon RD, Hodgson JA, Roy RR, Edgerton VR. (1999) Retention of hindlimb stepping ability in adult spinal cats after the cessation of step training. *Journal of Neurophysiology*, 81, 85-94.
- De Leon, RD, Hodgson JA, Roy RR, Edgerton VR. (1998) Locomotor capacity attributable to step training versus spontaneous recovery after spinalization in adult cats. *Journal of Neurophysiology*, 79, 1329-1340.
- Dobkin B. (1996). Recovery of locomotor control. *The Neurologist*, 2(4), 239-249
- Edgerton RE, Roy RR, Hodgson JA, Prober RJ, de Guzman CP, de Leon R. (1992) Potential of adult mammalian lumbosacral spinal cord to execute and acquire improved locomotion in the absence of supraspinal input. *Journal of Neurotrauma*, 9, S119-S128.
- Hodgson JA, Roy RR, de Leon, R, Dobkin B, Edgerton VR. (1994) Can the mammalian lumbar spinal cord learn a motor task? *Medicine and Science in Sports and Exercise*, 26, 1491-1497.
- Gruner JA, Altman J, Spivack N. (1980). Effects of arrested cerebellar development on locomotion in the rat. Cinematographic and electromyographic analysis. *Experimental Brain Research*, 40(4), 361-73.
- Lovely RG, Gregor RJ, Roy RR, Edgerton VR. (1990). Weight-bearing hindlimb stepping in treadmill-exercised adult spinal cats. *Brain Research*, 514, 206-218.
- Lovely RG, Gregor RJ, Roy RR, Edgerton VR. (1986) Effects of training on the recovery of full-weight-bearing stepping in the adult spinal cat. *Experimental Neurology*, 92, 421-435.

- Rossignol S, Drew T, Brustein E, Jiang W. (1999) Locomotor performance and adaptation after partial or complete spinal cord lesions in the cat. *Progress in Brain Research*, 123, 349-365.
- Smith JL, Edgerton VR, Eldred E, Zernicke RF. (1983) The chronic spinalized cat: A model for neuromuscular plasticity. *Birth defects original article series*, 19, 357-73.
- Suresh Babu R, Muthusamy R, Namasivayam A. (2000) Behavioral assessment of functional recovery after spinal cord hemisection in the bonnet monkey (*Macaca radiata*). *Journal of the neurological Sciences*, 178(2), 136-152.
- Whelan PJ, Pearson KG. (1997) Plasticity in reflex pathways controlling stepping in the cat. *Journal of Neurophysiology*, 78, 1643-1650.

Basic Science

4. MOTIVE CIRCUIT

- Armstrong D.M. (1986). Supraspinal contributions to the initiation and control of locomotion in the cat. *Progress in Neurobiology*, 26, 273-361.
- De Olmos JS, Heimer L. (1999) The concepts of the ventral striatopallidal system and extended amygdala. *Annals of the New York Academy of Sciences*, 877, 1-32.
- Groenewegen HJ, Wright CI, Beijer AV. (1996) The nucleus accumbens: gateway for limbic structures to reach the motor system? *Progress in Brain Research*. 107: p485-511.
- Ouchi Y, Okada H, Yoshikawa E, Nobezawa S, Futatsubashi M. (1999). Brain activation during maintenance of standing postures in humans. *Brain*, 122, 329-338.
- Shimamura M, Kogure I. (1983). Discharge patterns of reticulospinal neurons corresponding with quadrupedal leg movements in thalamic cats. *Brain Research*, 230, 27-34.

Basic Science

5. PHYSIOL/PHARM

- Adriani W, Felici A, Sargolini F, Roulet P, Usiello A, Oliverio A, Mele A. (1998) N-methyl-D-aspartate and dopamine receptor involvement in the modulation of locomotor activity and memory processes. *Experimental Brain Research*, 123(1-2), 52-9.
- Barbeau H, Rossignol S. (1990) The effects of serotonergic drugs on the locomotor pattern and on cutaneous reflexes of the adult chronic spinal cat. *Brain Research*, 514(1), 55-67.
- Barbeau H, Julien C, Rossignol S. (1987) The effects of clonidine and yohimbine on locomotion and cutaneous reflexes in the adult chronic spinal cat. *Brain Research*, 437(1), 83-96.
- Chau C, Barbeau H, Rossignol S. (1998) Effects of intrathecal alpha1- and alpha2-noradrenergic agonists and norepinephrine on locomotion in chronic spinal cats. *Journal of Neurophysiology*, 79(6): 2941-63.
- Chau C, Barbeau H, Rossignol S. (1998) Early locomotor training with clonidine in spinal cats. *Journal of Neurophysiology*, 79(1): 392-409.
- Giroux N, Brustein E, Chau C, Barbeau H, Reader TA, Rossignol S. (1998) Differential effects of the noradrenergic agonist clonidine on the locomotion of intact, partially and completely spinalized adult cats. *Annals of the New York Academy of Sciences*, 860, 517-20.
- Liste I, Rodriguez-Pallares J, Caruncho HJ, Labandeira-Garcia JL. (1999) Locomotor-activity-induced changes in striatal levels of preprotachykinin and preproenkephalin mRNA. Regulation by the dopaminergic and glutamatergic systems. *Brain Research. Molecular Brain Research*. 70(1): 74-83.
- Liste I, Guerra MJ, Caruncho HJ, Labandeira-Garcia JL. (1997) Treadmill running induces striatal Fos expression via NMDA glutamate and dopamine receptors. *Experimental Brain Research*, 115(3), 458-468.
- Marcoux J, Rossignol S. (2000) Initiating or blocking locomotion in spinal cats by applying noradrenergic drugs to restricted lumbar spinal segments. *Journal of Neuroscience*, 20(22) 8577-85.
- Remy-Neris O, Barbeau H, Daniel O, Boiteau F, Bussel B. (1999) Effects of intrathecal clonidine injection on

- spinal reflexes and human locomotion in incomplete paraplegic subjects. *Experimental Brain Research*, 129(3): 433-40.
- Rossignol S, Giroux N, Chau C, Marcoux J, Brustein E, Reader TA.(2001). Pharmacological aids to locomotor training after spinal injury in the cat. *Journal of Physiology*, 533(Pt 1), 65-74.
- Rossignol S, Chau C, Brustein E, Giroux N, Bouyer L, Barbeau H, Reader TA.(1998) Pharmacological activation and modulation of the central pattern generator for locomotion in the cat. *Annals of the New York Academy of Sciences*, 860, 346-59.
- Rossignol S, Barbeau H.(1993). Pharmacology of locomotion: an account of studies in spinal cats and spinal cord injured subjects. *The Journal of the American Paraplegia Society*, 16(4): 190-6.
- Pierce RC, Kalivas PW. (1997) A circuitry model of the expression of behavioral sensitization to amphetamine-like psychostimulants. *Brain Research. Brain Research Reviews*, 25(2), 192-216.

Clinical (Ailments)

1. SPINAL CORD INJURY

- Barrière G, Leblond H, Provencher J, and Serge Rossignol. (April 2008). Prominent Role of the Spinal Central Pattern Generator in the Recovery of Locomotion after Partial Spinal Cord Injuries. *The Journal of Neuroscience* 28(15): 3976-3987.
- Barbeau H, Fung J, Visintin M. (1999) New Approach to retrain gait in stroke and spinal cord injured subjects. *Neurorehabilitation and Neural Repair*, 13, 177-178.
- Barbeau H, Ladouceur M, Norman KE, Pepin A, Leroux A. (1999) Walking after spinal cord injury: Evaluation, treatment, and functional recovery. *Archives of Physical Medicine Rehabilitation*, 80, 225-235.
- Barbeau H, Rossignol S. (1994) Enhancement of locomotor recovery following spinal cord injury. *Current Opinion in Neurology*, 7, 517-524.
- Basso DM. (2000) Neuroanatomical substrates of functional recovery after experimental spinal cord injury: Implications of basic science for human spinal cord injury. *Physical Therapy*, 80, 808-817.
- † Behrman AL, et al. (2008). Locomotor Training Restores Walking in a Nonambulatory Child With Chronic, Severe, Incomplete Cervical Spinal Cord Injury *Physical Therapy* 88:580.
- † Behrman AL, Bowden MG, Nair PM. (2006) Neuroplasticity After Spinal Injury and Training: An Emerging Paradigm Shift in Rehabilitation and Walking Recovery. *Physical Therapy*, 86, 1406-1427.
- † Behrman AL, et al. (2005) Locomotor training progression and outcomes after incomplete spinal cord injury. *Physical Therapy*; 85:1356-1371.
- Behrman A, Harkema S. (2000) Locomotor Training After Human Spinal Cord Injury: A Series of Case Studies. *Phys The*, 80, 688-700
- † Bowden MG, et al. (2008) Beyond Gait Speed: A case report of multidimensional approach to locomotor rehabilitation outcomes in incomplete spinal cord injury. *Journal of Neurologic Physical Therapy*; 32:129-138.
- Bruehlmeier M, Dietz V, Leenders KL, Roelcke U, Missimer J, Curt A.(1998). How does the human brain deal with a spinal cord injury? *European Journal of Neuroscience*, 10(12), 3918-22.
- Calancie B, Lutton S, Broton.(1996) Central nervous system plasticity after spinal cord injury in man: Interlimb reflexes and the influence of cutaneous stimulation. *Electroencephalography and Clinical Neurophysiology*, 101, 304-315.
- Calancie B, Needham-Shropshire B, Jacobs P, Willer K, Zych G, Green BA. (1994) In voluntary stepping after chronic spinal cord injury: Evidenced for a central rhythm generator for locomotion in man. *Brain*, 117, 1143-1159.
- † Carvalho D Abreu DC, Junior AC, Rondina JM, Cendes F. (2008) Muscle hypertrophy in quadriplegics with combined electrical stimulation and body weight support training. *International Journal of Rehabilitation*

- Research, 31, 171-175.*
- † Carvalho D, Cliquet A. (2005) Energy expenditure during rest and treadmill gait training in quadriplegic subjects. *Spinal Cord* 43, 658–663
- † Carvalho D.C.L, Zanchetta M.C., Sereni J.M., Cliquet Jr A. (2005) Metabolic and cardiorespiratory responses of tetraplegic subjects during treadmill walking using neuromuscular electrical stimulation and partial body weight support. *Spinal Cord*, 43, 400-405.
- Colombo G, Wirz M, Dietz V.(2001). Driven gait orthosis for improvement of locomotor training in paraplegic patients. *Spinal Cord*, 39(5), 252-5.
- † Coupaud S, Jack LP, Hunt KJ, Allan DB, (2009)Muscle and bone adaptation after treadmill training in incomplete Spinal Cord Injury: a case study using peripheral Quantitative computed Tomography, *J Musculoskelet Neuronal Interact* 9(4): 288-297
- Dietz V, Nakazawa K, Wirz M, Erni T.(1999) Level of spinal cord lesion determines locomotor activity in spinal man. *Experimental Brain Research*, 128, 405-409.
- Dietz V, Wirz M, Colombo G, Curt A. (1998) Locomotor capacity and recovery of spinal cord function in paraplegic patients: A clinical and electrophysiological evaluation. *Electroencephalography and Clinical Neurophysiology*, 109, 140-153.
- Dietz V, Wirz M, Curt A, Colombo G. (1998). Locomotor pattern in paraplegic patients: Training effects and recovery of spinal cord function. *Spinal Cord*, 36, 380-390.
- Dietz V, Wirz M, Jensen L. (1997) Locomotion in Patients With Spinal Cord Injuries. *Physical Therapy*, 77(5),508-516
- Dietz V, Colombo G, Jensen L, Baumgartner L.(1995) Locomotor capacity of spinal cord in paraplegic patients. *Annals of Neurology*, 37, 574-582.
- † Dietz V, Colombo G, Jensen L.(1994) Locomotor activity in spinal man. *Lancet*, 344, 1260-1263.
- Dimitrijevic MR, Gerasimenko Y, Pinter MM. (1998) Evidence for a spinal central pattern generator in humans. *Annals of the New York Academy of Sciences*, 860, 360-376.
- † Ditor DS, MacDonald MJ, Kamath MV, Bugaresti J, Adams M, McCartney N, Hicks AL.(2005). The effects of body-weight supported treadmill training on cardiovascular regulation in individuals with motor-complete SCI. *Spinal Cord* 43, 664–673.
- † Dobkin B, et al (2006) Weight-Supported Treadmill vs Over Ground Training for Walking After Acute Incomplete SCI. *Neurology* 66: 484-493
- † Dobkin BH et al, (2003) Methods for a Randomized Trial of Weight-Supported Treadmill Training versus Conventional Training for Walking during Inpatient Rehabilitation after Incomplete Traumatic Spinal Cord Injury, *Neurorehabilitation and Neural Repair*, 17(3): 153-167
- Dobkin BH, Harkema S, Requejo P, Edgerton VR.(1995) Modulation of locomotor-like EMG activity in subjects with complete and incomplete spinal cord injury. *Journal of Neurologic Rehabilitation*, 9,183-190.
- Dobkin BH. (2000) Spinal and supraspinal plasticity after incomplete spinal cord injury: correlations between functional magnetic resonance imaging and engaged locomotor networks. *Progress in Brain Research*, 128, 99-111
- Dobkin B, Edgerton V, Fowler E. (1992) Sensory input during treadmill training alters rhythmic locomotor EMG output in subjects with complete spinal cord injury. *Proceedings of the Annual Meeting of the Society for Neuroscience*. Anaheim, CA,1403
- Dobkin B, Edgerton V, Fowler E, Hodgson J. (1992) Training induces rhythmic locomotor EMG patterns in subjects with complete SCI. *Neurology*, 42, 207-208.
- Dudley-Javoroski S, Shields R.K. (2008) Muscle and bone plasticity after spinal cord injury: Review of adaptations to disuse and to electrical muscle stimulation. *Journal of Rehab Research & Development*; 45,

283-296.

Edgerton VR, et al. (2001) Retraining the injured spinal cord. *Journal of Physiology*, 533.1, 15-22.

Field-Fote E, Dietz V. (2007) Single joint perturbation during gait: Preserved compensatory response pattern in spinal cord injured subjects. *Clinical Neurophysiology*, 118, 1607-1616.

† Field-Fote EC. (2001) Combined use of body weight support, functional electric stimulation, and treadmill training to improve walking ability in individuals with chronic incomplete spinal cord injury. *Archives of Physical Medicine and Rehabilitation*, 82(6), 818-24.

Field-Fote EC. (2000) Spinal cord control of movement: implications for locomotor rehabilitation following spinal cord injury. *Physical Therapy*, 80(5), 477-84.

† Giangregorio LM, Hicks AL, Webber CE, Phillips SM, Craven BC, Bugaresti JM and McCartney N. (2005) Body weight supported treadmill training in acute spinal cord injury: impact on muscle and bone. *Spinal Cord* 43, 649-657

† Gorassini MA, Norton JA, Nevett-Duchchere J, Roy FD, Yang JF. (2009) Changes in locomotor muscle activity after treadmill training in subject with incomplete spinal cord injury. *Journal of Neurophysiology*; 101 (2): 969-979

Harkema SJ, Hurley SL, Patel UK, Requejo PS, Dobkin BH, Edgerton VR. (1997) Human lumbosacral spinal cord interprets loading during stepping. *Journal of Neurophysiology*, 77, 797-811.

Herman R, D'Luzansky S, Willis W, Dilli S. (2002) Spinal cord stimulation facilitates functional walking in a chronic, incomplete spinal cord injured. *Spinal Cord*; 40, 65-68.

† Hicks AL, Martin Ginis KA. (2008) Treadmill training after spinal cord injury: It's not just about the walking. *Journal of Rehab Research & Development*; 45: 241-248.

† Hicks AL, et al. (2005) Long-term body-weight-supported treadmill training and subsequent follow-up in a person with chronic SCI: effects on functional walking ability and measures of subjective well-being. *Spinal Cord*, 43, 291-298

Hiersemenzel LP, Curt A, Dietz V. (2000) From spinal shock to spasticity: neuronal adaptations to a spinal cord injury. *Neurology*, 54(8), 1574-82.

Hulsebosch CE, Hains BC, Waldrep K, Young W. (2000) Bridging the gap: from discovery to clinical trials in spinal cord injury. *Journal of Neurotrauma*, 17(12), 1117-28.

Israel J, Campbell D, Kahn J, Hornby T. (2006) Metabolic Costs and Muscle Activity Patterns During Robotic- and Therapist-Assisted Treadmill Walking in Individuals with Incomplete Spinal Cord Injury. *Physical Therapy*, 86(11), 1466-1478.

† Knikou M, Angeli CA, Ferreira CK, Harkema SJ. (2008) Soleus H-reflex modulation during body weight support treadmill walking in spinal cord intact and injured subjects. *Experimental Brain Research*, 11p

Ladouceur M, Barbeau H. (2000) Functional electrical stimulation-assisted walking for persons with incomplete spinal injuries: changes in the kinematics and physiological cost of overground walking. *Scandinavian Journal of Rehabilitation Medicine*, 32(2), 72-9.

Ladouceur M, Pepin A, Norman KE, Barbeau H. (1997) Recovery of walking after spinal cord injury. *Advances in Neurology*, 72, 249-255.

Lajoie Y, Barbeau H, Hamelin M. (1999) Attentional requirements of walking in spinal cord injured patients compared to normal subjects. *Spinal Cord*, 37, 245-250.

Lynskey J.V., Belanger A, Jung R. (2008) Activity-dependent plasticity in spinal cord injury. *Journal of Rehab Research & Development*; 45: 229-240.

* Mayer R. (2007, October) Rocky road to recovery: Colorado's Craig Hospital specializes in TBI and SCI patients. *Advance for Physical Therapist and PT Assistants*, 10-11

Muir GD, Steeves JD. (1997) Sensorimotor stimulation to improve locomotor recovery after spinal cord injury.

- Trends in Neuroscience*, 20, 72-77.
- † Norman KE, Pepin A, Barbeau H. (1998) Effects of drugs on walking after spinal cord injury. *Spinal Cord*, 36(10), 699-715.
- † Norman KE, Pepin A, Ladouceur M, Barbeau H. (1995) A treadmill apparatus and harness support for evaluation and rehabilitation of gait. *Archives of Physical Medicine and Rehabilitation*, 76, 772-778.
- † Nymark J, DeForge D, Barbeau H, Badour M, Bercovitch S, Tomas J, Goudreau L, MacDonald J. (1998) Body weight support treadmill gait training in the subacute recovery phase of incomplete spinal cord injury. *Journal of Neurologic Rehabilitation*, 12, 119-138.
- Padula W, et al (2009) Modifying Postural Adaptation Following a CVA Through Prismatic Shift of Visuo-spatial egocenter. *Brain Injury*, 23(6): 566-576
- Pearson KG. (2000) Plasticity of neuronal networks in the spinal cord: modifications in response to altered sensory input. *Progress in Brain Research*, 128, 61-70.
- Pinter MM, Dimitrijevic MR. (1999) Gait after spinal cord injury and the central pattern generator for locomotion. *Spinal Cord*, 37, 531-537.
- * Prosser L. (2007) Locomotor training within inpatient rehabilitation program after pediatric incomplete spinal cord injury. *Physical Therapy*;87: 1224-1232
- † Protas EJ, Holmes SA, Qureshy H, Johnson A, Lee D, Sherwood AM. (2001) Supported treadmill ambulation training after spinal cord injury: a pilot study. *Archives of Physical Medicine and Rehabilitation*, 82(6), 825-31.
- † Puliti B. (March 2006) From Dreams to Destiny: Robotic body weight supported treadmill training teach SCI patients to walk again. *Advance for Physical Therapist and PT Assistants*, p42-43
- Requejo P, Harkema S, Edgerton V, Dobkin B. (1995) Direct relationship between the level of weight bearing and EMG activity during treadmill stepping in control and spinally injured subjects. *J Neurotrauma*, 12, 137
- Rossignol S, Chau C, Brustein E, Belanger M, Barbeau H, Drew T. (1996) Locomotor capacities after complete and partial lesions of the spinal cord. *Acta Neurobiologiae Experimentalis*, 56, 449-463.
- Wernig, A. (2006) Letter to the Editor- Long-term body-weight supported treadmill training and subsequent follow-up in persons with chronic SCI: effects on functional walking ability and measures of subjective well-being. *Spinal Cord*, 44, 265-266
- Wernig A, Nanassy A, Muller S. (1999) Laufband (treadmill) therapy in incomplete paraplegia and tetraplegia. *J Neurotrauma*, 16, 719-726.
- Wernig A. (1999) Laufband (treadmill) therapy in SCI persons. *Neurorehabilitation and Neural Repair*, 13, 175-176.
- Wernig A, Nanassy A, Muller S. (1998) Maintenance of locomotor abilities following Laufband (treadmill) therapy in para- and tetraplegic persons: Follow-up studies. *Spinal Cord*, 36, 744-749.
- Wernig A, Muller S, Nanassy A, Cagol E. (1995) Laufband therapy based on "Rules of Spinal Locomotion" is effective in spinal cord injured persons. *European Journal of Neuroscience*, 7, 823-829.
- Wernig A, Muller S. (1992) Laufband locomotion with body weight support improved walking in persons with severe spinal cord injuries. *Paraplegia*, 30, 229-238.
- † Wernig A, Muller S. (1991) Improvement of walking in spinal cord injured persons after treadmill training. In: Wernig A (ed.) *Plasticity of Motoneuronal Connections*. Amsterdam: Elsevier Science Publishers, 475-485
- † Whitman JM, et al. (2006) A comparison between two physical therapy treatment programs for patients with lumbar spinal stenosis: a randomized clinical trial. *Spine*, 31, 2541-2549.
- † Wickelgren I, et al. (1998) Teaching the Spinal Cord to Walk. *Science*, 279, 319-320.
- † Wirz M, Colombo G, Dietz V. (2001) Long term effects of locomotor training in spinal humans. *Journal of Neurology, Neurosurgery, and Psychiatry*, 71(1), 93-6.

Zomlefer M, Gaines R, McLeary L.(1983) Locomotor control in spinal cord injured humans. *Annual Meeting of the Society for Neuroscience*, 188 2p

Clinical (Ailments)

2. STROKE

- (2006) Study Helps Patients Walk Following Stroke. *Advance for Physical Therapist & PT Assistants*, p5
- Alexander M.(1997) A 36-year-old woman recuperating from a stroke. *JAMA*, 277(24), 1970-1976.
- † Barbeau H, Visintin M. (2003) Optimal outcomes obtained with body-weight support combined with treadmill training in stroke subjects. *Arch Phys Med Rehabilitation*, 84, 1458-65.
- † Breen J, Baker B, Snyder B, and Thibault K. (May 16, 2007) "Body Weight Support Treadmill Training Improves Walking in Sub-Acute and Chronic Severely Disabled Stroke Patients". *24th Annual Brain Injury & Stroke Conference Brain Injury Association of New Hampshire*.
- Byl NN, Pitsch EA, Abrams GM. (2008) Functional outcomes can vary by dose: Learning-based sensorimotor training for patients stable poststroke. *Neurorehabilitation and Neural Repair*,22(5), 494-504
- † Chen G, Patten C. (2006) Treadmill training with harness support: Selection of parameters for individuals with poststroke hemiparesis. *Journal of Rehabilitation Research & Development*, 43(4), 485-498
- † Chen G, Patten C, Kothari D, Zajac F. (2005) Gait deviations associated with post-stroke hemiparesis: improvement during treadmill walking using weight support, speed, stiffness, and handrail hold. *Gait and Posture*, 22, 57-62
- † Chong, D. (2007). Fall-Free Ambulation: Using body weight support gait training for a patient with pusher syndrome. *Advance for Physical Therapist & PT Assistants*, 18-19.
- † Combs S, Winchell Miller E, Forsyth E. (July 2007) Motor and functional outcomes of a patient post-stroke following combined activity and impairment level training. *Physiotherapy Theory and Practice*, 23
- † Daly J, et al. (2007) Intra-limb coordination deficit in stroke survivors and response to treatment. *Gait and Posture*, 25, 412-418
- † Danielsson A, Sunnerhagen KS.(2002). Oxygen consumption during treadmill walking with and without body weight support in patients with hemiparesis after stroke and in healthy subjects. *Archives of Physical Medicine and Rehabilitation*, 81, 953-957.
- † Davies PM. (1999) Weight-supported treadmill training. *Neurorehabilitation and Neural Repair*, 13, 167-169.
- Dean CM, Richard CL, Malouin F. (2000) Task-related circuit training improves performance of locomotor tasks in chronic stroke: A randomized, controlled pilot trial. *Archives of Physical Medicine and Rehabilitation*, 81, 409-417.
- Den Otter AR, Geurts A, Mulder T, Duysens J.(2006) Gait recover is not associated with changed in the temporal patterning of muscle activity during treadmill walking in patients with post-stroke hemiparesis. *Clinical Neurophysiology*,117, 4-15
- Dickstein R. (2008) Rehabilitation of gait speed after stroke: a critical review of intervention approaches. *Neurorehabilitation and Neural Repair*,22, 649-660
- Dobkin BH, (2000)The subacute stroke patient: Neurorehabilitation for stroke patients. In: Cohen, SN (ed) *Management of Ischemic Stroke*. McGraw-Hill, New York, 151-167.
- Dobkin BH, Gregor R, Rowler E. (1991) A strategy to train locomotion in patients with chronic hemiplegic stroke. *Annals of Neurology*, 30, 278.
- Eng J.J., Pang M.Y.C., Ashe M.C. (2008) Balance, falls, and bone health: Role of exercise in reducing fracture risk after stroke. *Journal of Rehab Research & Development*; 45(2):297-314.
- † Falmarin H, Golden S. (2008) Back on your feet: Balance programs help stroke survivors resume normal gait and activities. *Advance for Directors in Rehabilitation*,p49

- † Forrester LW, Wheaton LA, Luft AR. (2008) Exercise mediated locomotor recovery & lower-limb neuroplasticity after stroke. *Journal of Rehab Research & Development*, 45, 205-220.
- † Foster JB. (May 2007) Stroke rehabilitation study results surprise researchers. *Biomechanics*.
- Fritz S.L., Pittman A.L., Robinson A.C., Orton S.C., Rivers E.D. (2007) An Intense Intervention for Improving Gait, Balance, and Mobility for Individuals with Chronic Stroke: a pilot study. *Journal of Neurologic Physical Therapy*, 31, 71-76.
- Fritz SL, Light KE, Patterson TS, Behrman AL, Davis SB. (2005). Active Finger Extension Predicts Outcomes After Constraint-Induced Movement Therapy for Individuals With Hemiparesis After Stroke. *Stroke*; 36: 1172-1177
- Hafer-Macho CE, Ryan AS, Ivey FM, Macko RF. (2008) Skeletal muscle changes after hemiparetic stroke and potential beneficial effects of exercise intervention strategies. *Journal of Rehab Research & Development*; 45: 261-272
- Hase K, Fujiwara T, Tsuji T, Liu M. (2008) Effect of prosthetic gait training for stroke patients to induce use of paretic leg: a report of three cases. *Keio Journal of Med*, 57(3), 162-167
- Hassid E, Rose D, Commisaro J, Guttry M, Dobkin BH. (1997) Improved gait symmetry in hemiparetic stroke patients induced during body weight-supported treadmill stepping. *Journal of Neurologic Rehabilitation*, 11, 21-26.
- Hayes S, Carroll S. (1986). Early intervention care in the acute stroke patient. *Arch Phys Med Rehabil*, 67, 319-21
- Heinemann A, Roth E, Chichowski K, Betts H. (1987) Multivariate analysis of improvement and outcome following stroke rehabilitation. *Arch Neurol*, 44, 1167-1172
- † Hesse S, Sarkodie-Gyan T, Uhlenbrock D(1999). Development of an advanced mechanized gait trainer, controlling movement of the center of mass, for restoring gait in non-ambulant subjects. *Biomedizinische Technik*, 7(8), S194-201.
- Hesse S. (1999)Treadmill training with partial body weight support in hemiparetic patients – Further research needed. *Neurorehabilitation and Neural Repair*, 13, 179-181.
- † Hesse S, Uhlenbrock D, Sarkodie-Gyan T. (1999) Gait pattern of severely disable hemiparetic subjects on a new controlled gait trainer as compared to assisted treadmill walking with partial body weight support. *Clinical Rehabilitation*, 13, 401-410.
- † Hesse S, Konrad M, Uhlenbrock D. (1999) Treadmill walking with partial body weight support versus floor walking in hemiparetic subjects. *Archives of Physical Medicine and Rehabilitation.*; 80: 421-427.
- Hesse S, Jahnke MT, Schaffrin A, Lucke D, Reiter F, Konrad M.(1998) Immediate effects of therapeutic facilitation on the gait of hemiparetic patients as compared with walking with and without a cane. *Electroencephalography and Clinical Neurophysiology*, 109, 515-522.
- † Hesse S, Helm B, Krajnik J, Gregoric M, Mauritz KH. (1997) Treadmill training with partial body weight support: Influence of body weight release on the gait of hemiparetic patients. *Journal of Neurologic Rehabilitation*, 11, 15-20.
- † Hesse S, Malezic M, Schaffrin A, Mauritz K. (1995) Restoration of gait by combined treadmill training and multichannel electrical stimulation in non-ambulatory hemiparetic patients. *Scandinavian Journal of Rehabilitation Medicine*, 27, 199-204.
- † Hesse S, et al.(1995) Treadmill training with partial body weight support compared with physiotherapy in nonambulatory hemiparetic patients. *Stroke* 26(6), 976-981
- † Hesse S, Bertelt C, Schaffrin A, Malezic M, Mauritz KH.(1994) Restoration of gait in nonambulatory hemiparetic patients by treadmill training with partial body-weight support. *Archives of Physical Medicine and Rehabilitation*, 75, 1087-1093.

- † Hidler J, et al. (2009) Multicenter randomized clinical trial evaluating the effectiveness of the lokomat in subacute stroke. *Neurorehabilitation and Neural Repair*; 23: 5-13.
- Hornby GT, Campbell DD, Kahn JH, Demott T, Moore JL, Roth HR.(2008) Enhanced Gait-Related Improvements After Therapist- Versus Robotic-Assisted Locomotor Training in Subjects With Chronic Stroke. *Stroke*;39 :1786-1792
- Ivey F.M., Hafer-Macko C.E., Macko R.F. (2008) Task-oriented treadmill exercise training in chronic hemiparetic stroke. *Journal of Rehab Research & Development*; 45: 249-260.
- Jorgensen HS, Nakayama H, Raaschou HO, Olsen TS. (1995)Recovery of walking function in stroke patients: the Copenhagen Stroke Study. *Archives of Physical Medicine and Rehabilitation*, 76(1), 27-32.
- Kirker SGB, Simpson DS, Jenner JR, Wing AM. (2000) Stepping before standing: Hip muscle function in stepping and standing balance after stroke. *Journal of Neurology, Neurosurgery, and Psychiatry*, 68, 458-464.
- * Kline T, Schmit B, Kamper D.(2007) Exaggerated interlimb neural coupling following stroke. *Brain*, 159-169
- † Kosak, M.C, Reding, M.J. (2000) Comparison of Partial Body Weight-Supported Treadmill Gait Training Versus Aggressive Bracing Assisted Walking Post Stroke. *Neurorehabilitation and Neural Repair*, 14, 13-19
- † Laufer Y, Dickstein R, Chefez Y, Marcovitz E. (2001)The effect of treadmill training on the ambulation of stroke survivors in the early stages of rehabilitation: a randomized study. *Journal of Rehabilitation Research and Development*, 38(1), 69-78.
- Leonard, EMK. (1997)Treadmill versus floor ambulation in adults with hemiplegia: Changes in kinematic parameters. Master's Thesis. University of Wisconsin-Madison.
- Levine, P. (2008, January). Three Universal Truths. *Advanced for Directors in Rehabilitation*, 57.
- † Lewek D.M (2009) Allowing intralimb kinematic variability during locomotor training poststroke improves kinematic consistency: A subgroup analysis from a randomized clinical trial, *Physical Therapy* 89(8): 829-839
- Lindquist AR, Prado CL, Barros RM, Mattioli R, Lobo de Costa PH, Salvini TF. (2007) Gait training combining partial body-weight support, a treadmill, and functional electrical stimulation: effects on poststroke gait. *Physical Therapy*; 87: 1144-1154
- Lord J, Hall K. (1986) Neuromuscular reeducation vs. Traditional programs for stroke rehabilitation. *Arch Phys med Rehabil*, 67,89-91
- Logigian M, et al. (1983) Clinical exercise trial for stroke patients. *Arch Phys Med Rehabil*, 64, 364-367
- Macko RF, et al. (2008) Adaptive physical activity improves mobility function and quality of life in chronic hemiparesis. *Journal of Rehab Research & Development*; 45: 323-328.
- Malouin F, Potvin M, Prevost J, Richards C, Wood-Dauphinee S. (1992) Use of an intensive task-oriented gait training program in a series of patients with acute cerebrovascular accidents. *Physical Therapy* 72(11), 781-793
- † McCain KJ, Pollo FE, Baum BS, Coleman SC, Baker S, Smith PS. (2008) Locomotor treadmill training with partial body-weight support before overground gait in adults with acute stroke: a pilot study. *Arch Phys Med Rehabil*; 89: 684-691.
- McKenzie M.J., Yu S, Macko R.F., McLenithan J.C., Hafer-Macko C.E. (2008) Human genome comparison of paretic and nonparetic vastus lateralis muscle in patients with hemiparetic stroke. *Journal of Rehab Research & Development*; 45: 273-282.
- †* Miller, E., Combs, S.A., Fish, C., Bense, B., Owens, A., Burch A. (2008) Running Training After Stroke: A Single –Subject Report. *Physical Therapy*, 88, 1-12.
- †* Miller E.W., Quinn M.E., Seddon P.G. (2002) Body weight support treadmill and overground ambulation training for two patients with chronic disability secondary to stroke. *Physical Therapy*, 82(1), 53-61.
- Nadeau SE, et al. (2004) Donepezil as an adjuvant to constraint-induced therapy for upper-limb dysfunction after stroke: an exploratory randomized clinical trial. *Journal of Rehab Research and Development*; 41; 525-

- † Nilsson L, et al. (2001) Walking training of patients with hemiparesis at an early stage after stroke: a comparison of walking training on treadmill with body weight support and walking training on the ground. *Clinical Rehabilitation*, 15, 515-527
- Patterson S.L., Rodgers M.M., Macko R.F., Forrester L.W. (2008) Effect of treadmill exercise training on spatial and temporal gait parameters in subjects with chronic stroke: A preliminary report. *Journal of Rehab Research & Development*, 45, 221-228.
- Ploughman M, McCarthy J, Bosse M, Sullivan HJ, Corbett D. (2008) Does treadmill exercise improve performance of cognitive or upper-extremity tasks in people with chronic stroke? A randomized cross-over trial. *Archives of Physical Medicine and Rehabilitation*; 89: 2041-2047
- † Plummer P, Behrman A, Duncan P, Spigel P, Saracino D, Martin J, Fox E, Thigpen M, Kautz S. (2007) Effects of Stroke Severity and Training Duration on Locomotor Recovery After Stroke: A Pilot Study. *Neurorehabilitation and Neural Repair*, 21 (2): 137-151
- Pohl M, Werner C, Holzgraefe M, Kroczeck G, Wingendorg I, Hoolig G, Koch R, Hesse S. (2007) Repetitive locomotor training and physiotherapy improve walking and basic activities of daily living after stroke: a single-blind, randomized multicenter trial. *Clinical Rehabilitation*, 21. 17-27
- Puh U, Baer GD. (2008) A comparison of treadmill walking and overground walking in independently ambulant stroke patients: a pilot study. *Disability and Rehabilitation*: 1-9
- Resnick B, Michael K, Shaughnessy M, Kopunek S, Nahm ES, Macko RF. (2008) Motivators for treadmill exercise after stroke. *Topics in Stroke Rehabilitation*; 15 (5): 494-502
- Rimmer J.H., Wang E, Smith D. (2008) Barriers associated with exercise and community access for individuals with stroke. *Journal of Rehab Research & Development*; 45:315-322.
- † Segal, M (Aug/Sept. 2009) A Team Approach to Stroke Rehab. *Rehab Management*
- † Senior, R. (April 2008) Back On Track: Texas PT's research shows benefits of treadmill training following stroke. *Advance for Physical Therapist and PT Assistants*, p20
- Shepherd R, Carr J. (1999) Treadmill walking in neurorehabilitation. *Neurorehabilitation and Neural Repair*, 13, 171-173.
- Silver K, et al.(2000) Effects of Aerobic Treadmill Training on Gait Velocity, Cadent and Gait Symmetry in Chronic Hemiparetic Stroke: A Preliminary Report. *Neurorehabilitation and Neural Repair*, 14, 65-71
- Stein J. (2004) Motor Recovery Strategies After Stroke. *Top Stroke Rehabil*. 11, 12-22
- Stuart M, Chard S, Roettger S. (2008) Exercise for chronic stroke survivors: A policy perspective. *Journal of Rehab Research & Development*; 45: 329-336.
- † Sullivan K, et al (2007) Effects of task-specific locomotor and strength training in adult who were ambulatory after stroke: Results of the STEPS randomized clinical trial. *Physical Therapy*, 87, 1580-1602.
- † Sullivan KJ, Knowlton BJ, Dobkin B. (2002) Step Training with body weight support: effect of treadmill speed and practice paradigms on post stroke locomotor recovery. *Arch Phys Med Rehabil*, 83, 683-691.
- Suzuki K, et al. (1999) Relationship Between Stride Length and Walking Rate in Gait Training for Hemiparetic Stroke Patients. *American Journal of Physical Medicine & Rehabilitation*, 78, 147-152
- † Teixeira da Cunha Filho I, Lim PAC, Qureshy H, Henson H, Monga T, Protas E. (2002) Gait outcomes after acute stroke rehabilitation with supported treadmill ambulation training: a randomized controlled pilot study. *Arch Phys Med Rehabil*, 83, 1258-1265
- † Teixeira da Cunha Filho I, Lim PA, Qureshy H, Henson H, Monga T, Protas EJ.(2001) A comparison of regular rehabilitation and regular rehabilitation with supported treadmill ambulation training for acute stroke patients. *Journal of Rehabilitation Research and Development*, 38(2), 245-55.
- Tilson JK, Settle SM, Sullivan KJ. (2008) Application of evidence-based practice strategies: current trend in

walking recover interventions poststroke. *Topics in stroke rehabilitations*;15 (3): 227-246

- †* Trueblood P. (2001) Partial body weight treadmill training in persons with chronic stroke. *NeuroRehabilitation*, 16, 141-153
- † Visintin M, Barbeau H, Korner-Bitensky N, Mayo NE. (1998) A new approach to retrain gait in stroke patients through body weight support and treadmill stimulation. *Stroke*, 29, 1122-1128.
- † Visintin M, Barbeau H. (1994) The effect of parallel bars, body weight support and speed on the modulation of the locomotor pattern of spastic paretic gait. A preliminary communication. *Paraplegia*, 32, 540-553.
- † Visintin M, Barbeau H.(1989) The effect of body weight support on the locomotor pattern of spastic paretic patients. *Canadian Journal of Neurological Science*, 16, 315-325.
- † Werner C, Bardeleben A, Mauritz K-H, Kirker S, Hesse S. (2002). Treadmill training with partial body weight support and physiotherapy in stroke patients: a preliminary comparison. *European Journal of Neurology*, 9, 639-644
- † Werner C, Von Frankenberg S, Treig T, Konrad M, Hesse S. (2002) Treadmill training with partial body weight support and an electromechanical gait trainer for restoration of gait in subacute stroke patients: a randomized cross over study. *Stroke*, 33, 2895-2901

Clinical (Ailments)

3. CEREBRAL PALSY

- †* Begnoche, DM, Pitetti, K. (2007) Effects of Traditional Treatment and Partial Body Weight Treadmill Training on the Motor Skills of Children with Spastics Cerebral Palsy a Pilot Study. *Pediatric Physical Therapy*, 19, 11-19.
- † Borggraefe I, et al (2008) Improved Gait Parameters After Robotic-Assisted Locomotor Treadmill Therapy in a 6-Year-Old Child with Cerebral Palsy. *Movement Disorder*, 23(2): 280-83
- † Bundonis J. (2003, October) Up and Walking- A case example of partial weight bearing treadmill training. *Advance for Physical Therapists & PT Assistants* 13(23), 39-40
- †* Cherg R-J, Liu C-F, Lau T-W, Hong R-B. (2007) Effect of treadmill training with body weight support on gait and gross motor function in children with spastic cerebral palsy. *Am J Phys Med Rehabil*, 86, 548-555.
- Day J, et al. (2004) Locomotor Training with Partial Body Weight Support on a Treadmill in a Nonambulatory Child with Spastic Tetraplegic Cerebral Palsy: A Case Report. *Pediatric Physical Therapy*, 16, 106-113.
- †* Dodd KJ, Foley S. (Feb 2007) Partial body-weight-supported treadmill training can improve walking in children with cerebral palsy: a clinical controlled trial. *Dev Med Child Neurol* 49(2): 101-5
- †* Flanagan N, Jacoby N, Johnson P, Krzak. Short Term Intensive: Physical Therapy to Address Balance and Gait Efficiency in an Adolescent who has Cerebral Palsy.
- †* Krzak J, Flanagan A, Johnson P, Jacoby. Preliminary Study of the Effects of Short Term Intensive treadmill training for children who have Cerebral Palsy.
- Leonard CT, Hirschfeld H, Forssberg H.(1991) The development of independent walking in children with cerebral palsy. *Developmental Medicine and Child Neurology* 33(7), 567-77.
- † Mattern-Baxter K. (2009) Effect of partial body weight supported treadmill training on children with cerebral palsy. *Pediatric Physical Therapy*, 21, 12-22
- † McNevin NH, Coraci L, Schafer J.(2000) Gait in adolescent cerebral palsy: The effect of partial unweighting. *Archives of Physical Medicine and Rehabilitation*, 81, 525-8.
- † Phillips J, et al. (2007) Ankle dorsiflexion fMRI in children with cerebral palsy undergoing intensive body-weight-supported treadmill training: a pilot study. *Developmental Medicine & Children Neurology*,49, 39-44
- †* Provost B, et al. (2007) Endurance and Gait in Children With Cerebral Palsy After Intensive Body Weight-Supported Treadmill Training. *Pediatric Physical Therapy*.
- † Richards CL, Malouin F, Dumas F, Marcoux S, Lepage C, Menier C. (1997) Early and intensive treadmill

locomotor training for young children with cerebral palsy: A feasibility study. *Pediatric Physical Therapy*, 9, 158-65

Russell SD, Bennett BC, Kerrigan DC, Abel MF. (2007) Determinants of gait as applied to children with cerebral palsy. *Gait and Posture*, 26, 295-300

† Schindl MR, Forstner C, Kern H, Hesse S.(2000) Treadmill training with partial body weight support in nonambulatory patients with cerebral palsy. *Archives of Physical Medicine and Rehabilitation*, 81, 301-306.

Trahan J, Malouin F.(2002) Intermittent intensive physiotherapy in children with cerebral palsy: a pilot study. *Developmental Medicine & Child Neurology*, 44, 233-239

Clinical (Ailments)

4. HEAD INJURY

†* Brown T, Mount J, Rouland BL, Kautz KA, Barnes RM, Kim J.(2005) Body Weight-Supported Treadmill Training Versus Conventional Gait Training for People with Chronic Traumatic Brain Injury. *J Head Trauma Rehabil*,20(5), 402-415

†* Moriello, G. Frear, M., Seaburg K. (2009) The Recovery of Running Ability in an Adolescent Male After Traumatic Brain Injury: A Case Study. *Journal of Neurologic Physical Therapy*, 33 (2), 111-120

†* Mossberg K, Orlander E, and Norcross J. (2008) Cardiorespiratory capacity after weight-supported treadmill training in patients with traumatic brain injury. *Physical Therapy*, 87, 1-11.

† Scherer M. (2007) Gait rehabilitation with body weight-supported treadmill training for a blast injury survivor with traumatic brain injury. *Brain Injury*; 21,93-100

†* Seif-Naraghi AH, Herman RM. (1999)A novel method for locomotion training. *Journal of Head Trauma Rehabilitation*, 14, 146-162.

Wainberg M, Barbeau H. (1982) Applicability of progressive weight bearing in rehabilitation of neurologically impaired gait. *Can J Neurol Sci*, 12.

† Wilson DJ, Powell M, Gorham JL, Childers M (2006) Ambulation training with and without partial weightbearing after traumatic brain injury: Results of a randomized, controlled trial. *American Journal of Physical Medicine & Rehabilitation*; 85:68–74

†* Wing, K. (December 2007) Living Proof: Mobility Research offers neurological patient light at the end of the tunnel. *Advance for Directors in Rehabilitation*, p47

Clinical (Ailments)

5. PARKINSON'S

Bello O, Sanchez JA, Fernandez-del-Olmo M. (2008) Treadmill walking in Parkinson's Disease patients: adaptation and generalization effect. *Movement Disorders*;23(9) 1243-1249.

Broughton, B. (2004, April) Parkinson's researchers take steps toward improving motor deficits. *Biomechanics*, 43

Dietz V, Colombo G.(1998) Influence of body load on the gait pattern in Parkinson's disease. *Movement Disorders*, 13, 255-261.

† Fisher BE, et al. (2008). The effect of exercise training in improving motor performance and corticomotor excitability in people with early Parkinson's Disease. *Archive of Physical Medicine and Rehabilitation*, 89, 1221-1229

† Fisher B, Fong M, Yip J.(2004, August) Treadmill training improves function in Parkinson's patients. *Biomechanics*, 57-61

Hanankawa T, Fukuyama H, Katsumi Y, Honda M, Shibasaki H.(1999) Enhanced lateral premotor activity during paradoxical gait in Parkinson's disease. *Annals of Neurology*, 45, 329-336.

Hanankawa T, Katsumi Y, Fukuyama H, Honda M, Hayashi T, Kimura J, Shibasaki H.(1999) Mechanisms

underlying gait disturbance in Parkinson's disease: A single photon emission computed tomography study. *Brain*, 122, 1271-1282.

- † Miyai I.(2002) Long-Term Effect of Body Weight-Supported Treadmill Training in Parkinson's Disease: A Randomized Controlled Trial. *Arch Phys Med Rehabil*, 83, 1370-3
- † Miyai I, Fujimoto Y, Ueda Y, Yamamoto H, Nozaki S, Saito T, Kang J. (2000) Treadmill training with body weight support: Its effect on Parkinson's disease. *Archives of Physical Medicine and Rehabilitation*. 81, 849-852.
- Morris M, Matyas T, Iansek R, Summers J. (1996) Temporal stability of gait in parkinson's disease. *J Phys Ther* 76(7), 763-777
- Pahapill PA, Lozano AM. (2000) The pedunculopontine nucleus and Parkinson's disease. *Brain*, 123 (Pt 9), 1767-83.
- † Toole T, Maitland C.G., Warren E, Hubmann M.F., Panton L. (2005) The effects of loading and unloading treadmill walking on balance, gait, fall risk, and daily function in Parkinsonism. *NeuroRehabilitation*, 20, 307-322.

Clinical (Ailments)

6. OTHER

- Barbeau H, Fung J. (1992) New experimental approaches in the treatment of spastic gait disorders. *Med Sport Sci*, 36, 234-246
- † Barbeau H, Blunt R. (1991) A Novel interactive locomotor approach using body weight support to retrain gait in spastic paretic subjects. In: Wernig A (ed.). *Plasticity of Motoneuronal Connections*. Amsterdam: Elvsevier Science Publishers,461-474
- † Beer S, et al. (2008) Robot-assisted gait training in multiple sclerosis: a pilot randomized trial. *Multiple Sclerosis*, 14, 231-236
- † Hesse S, et al. (2003) Treadmill Training With Partial Body-Weight Support After Total Hip Arthroplasty: A Randomized Controlled Trial. *Arch Phys Med Rehabil*, 84, 1767-73
- Hunter D, Smith Cole E, Murray JM, Murray TD. (1995) Energy expenditure of below-knee amputees during harness-supported treadmill ambulation. *The Journal of Orthopaedic and Sports Physical Therapy*, 21(5), 268-76.
- Kanai A, Kiyama T, Genda E, Suzuki Y. (2008) Biomechanical investigation of ambulatory training in patients with acetabula dysplasia. *Gait and Posture*; 28: 52-57
- * Kline Mangione K, Axen K, Haas F. (1996) Mechanical unweighting effects on treadmill exercise and pain in elderly people with osteoarthritis of the knee. *Physical Therapy*, 76, 387-394.
- Liston R, Mickelborough J, Harris B, Hann A.W., Tallis R.C. (2000) Conventional physiotherapy and treadmill re-training for higher-level gait disorders in cerebrovascular disease. *Age and Ageing*, 29, 311-318.
- † Lo AC, Triche EW. (2008) Improving gait in Multiple Sclerosis using robot-assisted, body weight supported treadmill training. *Neurorehabilitation and Neural Repair*; 22: 661-671
- Svendsen B. (1996, March) Treatment of the hemiplegic patient: new strategies for gait training. *Phys Ther Prod*, 32-34
- Tuckey J, Greenwood. (2004). Rehabilitation after severe Guillain-Barre syndrome: the use of the partial body weight support. *Physiotherapy Research International*; 9: 96-103.
- Waagfjord J, Levange P, Certo C. (1990) Effects of Treadmill training on gait in a hemiparetic patient. *Phys Ther* 70, 549-560
- Werner C., Lindquist AR., Bardeleben A, Hesse S. (2007) The Influence of Treadmill Inclination on the Gait of Ambulatory Hemiparetic Subjects. *Neurorehabilitation and Neural Repair* 21: 76-80.

Clinical

PEDIATRICS

- Angulo-Barroso R, Burghardt AR, Lloyd M, Ulrich DA. (2008) Physical activity in infants with Down syndrome receiving a treadmill intervention. *Infant Behav Dev* 31(2): 255-69.
- Bly L. (1994) Motor Skills Acquisition in the First Year. Tucson, Arizona: *Therapy Skill Builders*.
- Campbell S, et al. (1999) *Physical Therapy for Children*. Philadelphia: Churchill Livingstone.
- †* Cernak K, Stevens V, Price R, Shumway-Cook A. (2008) Locomotor training using body-weight support on a treadmill in conjunction with ongoing physical therapy in a child with severe cerebella ataxia. *Physical Therapy*, 88, 88-97.
- † Damiano DL, DeJong SL (2009) A systematic review of the effectiveness of treadmill training and body weight support in pediatric rehabilitation. *Neurological Physical Therapy*, 33(1), p27
- Kluger J (2007/April) Walking Away from Paralysis. *Time*
- * Mattern- Baxter, K , Bellamy, S, Mansoor, J Effects of Intensive Locomotor Treadmill Training on Young Children with Cerebral Palsy, *Pediatric Physical Therapy* 308-319
- Musselman KE and Yang, JF. (2007) Loading the Limb During Rhythmic Leg Movements Lengthens the Duration of Both Flexion and Extension in Human Infants. *J Neurophysiol* 97: 1247-1257
- †* Paleg, G. (2007, November). When Needs are Special: Determining mobility requirements for a child with Joubert syndrome. *Rehab Pub*, 24-28.
- * Paleg G. (2005, April) Standing Strong: An update on passive standing programs for children. *Advance for Physical Therapist and PT Assistants*, 39
- †* Paleg G. (2002, March) Walking the Walk: One therapist details her techniques to help children facilitate walking using partial weight bearing treadmill training. *Advance for Physical Therapists & PT Assistants*, 41-42
- †* Puliti, B (March 2006). Fighting for Their Future: Stony Brook University Medical Center sets standard in pediatric MS care. *Advance for Physical Therapist and PT Assistants*, p36-37
- † Santoyo M (2009) Reaching Past Spina Bifida – Early PT Enhances Future Function Today In PT
- Schnidehette S, Breu G. (2002) One Giant Leap. *People*, p 82-84
- Sutherland D. (1997) The Development of mature gait. *Gait & Posture*, 6, 163-170
- Ulrich DA, Lloyd MC, Tiernan CW, Looper JE, Angulo-Barroso RM. (January 2008). Effects of Intensity of Treadmill Training on Developmental Outcomes and Stepping in Infants With Down Syndrome: A Randomized Trial. *Physical Therapy* 88: 114-122
- Ulrich, DA & Ulrich, BSD.(1999) Treadmill training facilitates the onset of walking in infants with Down Syndrome. Research review from the center for Motor Behavior, University of Michigan
- Woo R. (2001, January) Spasticity: orthopedic perspective. *Journal Child Neurology*, 16(1),47:53
- Yang JF, Stephens MJ, Vishram R.(1998, March) Infant stepping: a method to study the sensory control of human walking. *J Physiol*,507 (pt 3),927-937

Clinical

MISCELLANEOUS

- Barbeau H, McCrea DA, O'Donovan MJ, Rossignol S, Grill WM, Lemay MA. (1999) Tapping into spinal circuits to restore motor function. *Brain Research Reviews*, 30, 27-51.
- Barbeau H, Norman K, Fung J, Visintin M, Ladouceur M. (1998) Does neurorehabilitation play a role in the recovery of walking in neurological populations? *Annals of the New York Academy of Sciences*, 860, 377-92.
- Barker S.P., Freedman W, Hillstrom H. (2006) A novel method of producing a repetitive dynamic signal to examine reliability and validity of gait analysis systems. *Gait and Posture*, 24, 448-452.
- Bastiaanse CM, Duysens J, Dietz V. (2000) Modulation of cutaneous reflexes by load receptor input during human walking. *Experimental Brain Research*, 135(2), 189-98.

- Boylstein C, Rittman M, Gubrium J, Behrman A, Davis, S. (2005). The social organization in constraint-induced movement therapy. *Journal of Rehabilitation Research & Development*; 42: 263-276.
- Capaday C, Lavoie BA, Barbeau H, Schneider C, Bonnard M. (1999) Studies on the corticospinal control of human walking. I. Responses to focal transcranial magnetic stimulation of the motor cortex. *Journal of Neurophysiology*, 81, 129-139.
- Conroy EB, (2009) Hospital based stroke rehabilitation in the United States, *Topics in Stroke Rehabilitation* 16(1): 34-43
- Cooper RA, et al. (1999). Gait Analysis in Rehabilitation Medicine. *American Journal of Physical Medicine & Rehabilitation*, 78, 278-280.
- † Crompton S, et al. (2001) Practical issues in retraining walking in severely disabled patients using treadmill and harness support systems, *Australian Journal of Physiotherapy* 47() 211-213
- Davis BL, Cavanagh PR. (1993) Simulating reduced gravity: a review of biomechanical issues pertaining to human locomotion. *Aviation, Space, and Environmental Medicine*, 64(6), 557-66.
- Faist M, Blahak C, Duysens J, Berger W. (1999) Modulation of the biceps femoris tendon jerk reflex during human locomotion. *Experimental Brain Research*, 125(3), 265-70.
- Field-Fote EC, Hufford T, Kerfoot N, Kizer S, Calancie B. (2000). Effect of lower extremity weight bearing load on motorneuron excitability in able-bodied subjects. *Electromyography and Clinical Neurophysiology*, 40(8), 459-64.
- Finch L, Barbeau H. (1986) Hemiplegic gait: new treatment strategies. *Physiotherapist Canada*, 38, 36-41.
- † Fiske J. (2005). Support System: Application for bodyweight-supported treadmill training stretch beyond post-stroke and spinal cord patients. *Advance for Directors in Rehabilitation*, 45
- Forsberg H, Hirschfeld H. (1988) Phasic modulation of postural activation patterns during human walking. *Progress in Brain Research*, 76, 221-227.
- Fox EL, Bartels RL, Chaloupka EC, Klinzing JE, Hoche J. (1975) Oxygen cost during exercise in simulated subgravity environments. *Aviation, Space, and Environmental Medicine*, 46(3), 300-3.
- Friedman P, Richmond D, Baskett J. (1988). A prospective trial of serial gait speed as a measure of rehabilitation in the elderly. *Age and Ageing*, 17, 227-235
- † Giesser B, et al. (2007) Locomotor training using body weight support on a treadmill improves mobility in person with multiple sclerosis: a pilot study. *Multiple Sclerosis*, 13, 224-231.
- Hellstrad E. (1985) Brain metabolism and human gait – a PET study. *Acta Physiology Scandinavia*, 125, 555-556.
- Huddleston OL (1961). *Therapeutic Exercises; Kinesiotherapy*. Philadelphia, FA Davis Co.
- Light KJ (2008) People, not robots, prove to be better walking assistants. *In Stride Gait Rehab*. September 2008, 2p
- Morrison S, Backus D. (2007). Locomotor Training: Is Translating Evidence Into Practice Financially Feasible? *JNPT*, 31, 50-54
- † Pillar T, Dickenstein R, Smolinski Z. (1991) Walking Re-education with Partial Relief of Bodyweight Support in Rehabilitation of Patients with Locomotor Disabilities. *Journal of Rehabilitation Research*, 28, 47-52.
- Riley PO, Paolini G, Della Croce U, Paylo KW, Kerrigan DC. (2007) A kinematic and kinetic comparison of overground and treadmill walking in healthy subjects. *Gait & Posture*, 26, 17-24
- Scheets P, Sahrman S, Norton B. (2007) Use of movement system diagnoses in the management of patients with neuromuscular conditions: a multiple-patient case report. *Physical Therapy*, 87
- Serino A, Barbiani M, Rinaldesi ML, Ladavas E (2009) Effectiveness of prism adaptation in neglect rehabilitation: a controlled trial study. *Stroke*, 40, 1392-1398
- Sinkjaer T, Andersen JB, Ladouceur M, Christensen LO, Nielsen JB. (2000) Major role for sensory feedback in

soleus EMG activity in the stance phase of walking in man. *Journal of Physiology*, 523, 817-827.

Stephens JM, Yang JF.(1999) Loading during the stance phase of walking in humans increases the extensor EMG amplitude but does not change the duration of the step cycle. *Experimental Brain Research*, 124, 363-370.

† Studer, M. (June 2008) Keep it Moving: Advances in gait training techniques help clients reduce balance issues and limit their incidence of falls. *Rehab Management*, 7p

Van de Crommert HWAA, Mulder T, Duysens J. (1998) Neural control of locomotion; Part 2: Sensory control of the central pattern generator and its relation to treadmill training. *Gait and Posture*, 7, 251-263.

Van Faassen F, Molen NH.(1973) A new method in the treatment of gait disorders. *Medicine and Sport*, 8, 489-491.

Wainberg M, Barbeau H. (1985) Applicability of progressive weight bearing in rehabilitation of neurologically impaired gait. *Le Journal Canadien des Sciences Neurologiques*, 12, 183.

Zehr PE, Stein RB.(1999) What functions do reflexes serve during human locomotion? *Progress in Neurobiology*, 58, 185-205.

Clinical

GENERAL BWS-GT

† Daly J, Ruff R. (2004) Feasibility of combining multi-channel functional neuromuscular stimulation with weight-supported treadmill training. *Journal of the Neurological Sciences*, 255, 105-115.

† Dietz V (2008) Body weight supported gait training: From laboratory to clinical setting *Brain Research Bulletin*, 76: 459-463

† Dobkin BH. (1999) An overview of treadmill locomotor training with partial body weight support: A neurophysiologically sound approach whose time has come for randomized clinical trials. *Neurorehabilitation and Neural Repair*, 13, 157-165.

† Franz J, et al. (2007) Physiological modulation of gait variable by an active partial body weight support system. *Journal of Biomechanics*, 40, 3244 -3250

† Finch L, Barbeau H, Arsenault B. (1991) Influence of body weight support on normal human gait: development of a gait retraining strategy. *Phy Ther*, 71(11).

† Finch L, Barbeau H. (1985). Influence of partial weight bearing on normal human gait: the development of gait retraining strategies. *Can J Neurol Sci* 12,183.

† Hesse S. (2001) Locomotor Therapy in NeuroRehabilitation. *NeuroRehabilitation*, 16, 133-139.
Koch P. Integrated Therapy to Improve Walking.

† Mauritz K. (2002) Gait training in hemiplegia. *European Journal of Neurology*, 9, 23-29

† O'Rourke J (2009/April) Counterpoise: Getting up and around, safely, after balance deficit. *Rehab Management*, 22-24

† Pollock R, Heneghan P, Riches PE. (2008) Under-arm partial body weight unloading causes spinal elongation and vibration attenuation during treadmill walking. *Gait & Posture*; 28: 271-277

* † Seif-Naraghi A.(2003). Walking-Spinal or Cortical? *NDTA Network*,9

† Selzer ME. (1999) Partial body weight-supported treadmill training in neurorehabilitation: Has the case been made? *Neurorehabilitation and Neural Repair*, 13, 155.

† Studer M. (2007) Invited Commentary: Giving Your Outpatient Rehabilitation Program a Lift- The financial Feasibility of Body Weight Support Treadmill Training. *JNPT*, 31, 54-55

† Wesenhagen S (2008) Positive Steps: A case report on using BWSTT combined with over ground ambulation. *Advance for Physical Therapist and PT Assistants*. 20-21

† Wilson D (2007) Principles of Gait Rehabilitation and the Efficacy of Partial Body-Weight-Supported Training. *Critical Reviews™ in Physical and Rehabilitation Medicine* 19: 169-194

* † Wolski C. (October 2005). Real-World Rehab– Technological Locomotion. *Rehab Management*, p22

Clinical DEVICES

- † Aaslund MK, Moe-Nilssen R. (2008) Treadmill walking with body weight support. Effect of treadmill, harness and body weight support systems. *Gait and Posture*, 28, 303-308
- Barbeau H, Wainberg M, Finch L.(1987) Description and application of a system for locomotor rehabilitation. *Medical and Biological Engineering and Computing*, 25, 341-344.
- Bundonis J. (2005, July) Walking Toward the Goal - When walkers and gait trainers do not offer enough assistance for gait training what else can be offered? *Advance for Physical Therapists & PT Assistants*, 16(15), 47
- * Cunningham, M. (2005, April) Stepping Forward: Mobility Research's LiteGait System adapts to subacute and outpatient care. *Advance for Directors in Rehabilitation*, 62.
- † Farrimond K.(1989) Parachute Harness. *Physiotherapy*, 75, 740-741.
- † Galbreath R. (2004, January/February) Worth the Weight. *Physical Therapy Products*, 34-36
- † Gazzani F, Bernardi M, Macaluso A, Coratella D, Ditunno JF Jr, Castellano V, Torre M, Macellari V, Marchetti M.(1999) Ambulation training of neurological patients on the treadmill with a new Walking Assistance and Rehabilitation Device (WARD). *Spinal Cord*, 37(5), 336-44.
- * Good, K. (2003, October) On the Move: LiteGait offer a novel setting for mobility training. *Advance for Directors in Rehabilitation*, 92.
- Harburn KL, Hill KM, Kramer JF, Noh S, Vandervoort AA, Matheson JE.(1993) An overhead harness and trolley system for balance and ambulation assessment and training. *Archives of Physical Medicine and Rehabilitation*, 74(2), 220-3.
- †* Hastings S. (2002, January) A Single Step. Mobility Research's partial weight-bearing treadmills help patients gain functions. *Advance for Directors in Rehabilitation*, 64
- Hollis M, Roper MHS. (1958). *Suspension Therapy in Rehabilitation*. Baltimore, Williams and Wilkins Co.
- Hosalkar, HS; Reddy, S; Keenan, MA. (2007) What's New in Orthopaedic Rehabilitation. *The Journal of Bone and Joint Surgery* 89:2316-2324.
- Kawamura J, Ide T, Hayashi S, Ono H, Honda T.(1993) Automatic suspension device for gait training. *Prosthetics and Orthotics International*, 17, 120-125.
- †* Koch P. (2004, July) Reaching Milestones: The Walkable helps pediatric clients achieve mobility goals. *Advance for Directors in Rehabilitation*, 76.
- Newman DJ, Alexander HL, Webbon BW. (1994) Energetics and mechanics for partial gravity locomotion. *Aviation, Space, and Environmental Medicine*, 65(9), 815-23.
- Paleg G. (2004) Gateway to Gait: Partial weight bearing treadmill training and gait trainer can assist clients with all level of gross motor dysfunction. *Rehab Management*, 28-31
- * Sisto SA, Forrest GF, Faghiri PD. (2008) Technology for Mobility and Quality of Life in Spinal Cord Injury. *IEEE Engineering in Medicine and Biology Magazine*, 52-68.
- Studer M, Whetten B (2008/December) Customer Support: Mobility Research's LiteGait aids patients and clinicians. *Advance for Directors in Rehabilitation*, p41
- Wilson MS, Qureshy H, Protas EJ, Holmes SA, Krouskop TA, Sherwood AM.(2000) Equipment specifications for supported treadmill ambulation training. *Journal of Rehabilitation Research and Development*, 37(4) 415-22.